
# **COOPDECH Endobronchial Blocker Tube** • Product Composition

Туре	Product Number	Bronchial Blocker Tube	Auto-inflator	Joint Connector	Cuff	Number of Items Included (set)
Type A	BBT-A3060-EU	Outer diameter 3.0mm/	_	0	Rectangular round-	1
Type B	BBT-B3060-EU	Length 600mm	0	0	shaped	1
Type A	BBT-A3060S-EU	Outer diameter 3.0mm/	_	0	Small spindle-	1
Type B	BBT-B3060S-EU	Length 600mm	0	0	shaped	1

#### 1set/box Sterilized

• Please note that specifications and outward appearance may change without notice for improvement purposes.



#### PRECAUTION FOR USE

- During the intubation of the endobronchial blocker, sufficiently exercise caution against the risk of tracheal or bronchial injury while monitoring using the fiberoptic bronchoscope.
- The endobronchial blocker tube should not be advanced if resistance is encountered.
- The markings observed in the endobronchial blocker are a rough standard to monitor the depth of intubation. The physician has to clinically determine the actual depth of intubation.
- The intubation site of the endobronchial blocker tube should be regularly monitored by fiberoptic bronchoscope.
- When the posture of the patient is changed (postural change is observed), never fail to check the location of intubated endobronchial blocker tube.
- Do not use a laser device or electrical surgical device near the endobronchial blocker tube
- If the physician notices difficulty in ventilation during bronchial blockage, deflate the cuff promptly, check the condition of the patient and take appropriate actions.
- Keep watching and make sure that the patient is under normal status.
   Once it is judged that the abnormality is occurred due to the product, take appropriate actions immediately to make sure the patient safety.

- Do not use oil-based lubricants such as Paraffin oil, Olive oil and Glycerin because they may cause deterioration of the cuff.
- 10. Do not use lidocaine aerosols for topical application because they may erase the markings
- 11. Do not apply the lubricant to the edge of the endobronchial blocker tube because it may cause occlusion of the cavity of the endobronchial blocker tube
- 12. Check if the endotracheal tube connector and the ventilation port of the joint connector are firmly connected with the endotracheal tube and the anesthetic route, respectively.
- 13. Always monitor the status of patient by measuring the oxygen saturation using pulse oximeter, the changing of intrinsic airway pressure and expiratory carbon dioxide.
- 14. The right upper lobe may be occluded if the right main bronchus is blocked
- 15. Do not use the device in an ozone-rich atmosphere.

Pioneering the future of medical society

DAIKEN MEDICAL CO., LTD.

International Department: 2-6-2, Ayumino, Izumi-city, Osaka, 594-1157, Japan

COOPDECH's product information is available at the corporate website. http://www.daiken-iki.co.jp/ [Agency]

# 2020.12 Contents of this catalog are as of December 2020.


# E COOPDECH

# **COOPDECH Endobronchial Blocker Tube**

- COOPDECH Endobronchial Blocker Tube allows one-lung ventilation using a conventional endotracheal tube and a fiber optics bronchoscope.
- Upon completion of one-lung ventilation, there is no need to replace trachea tubes, deflate the balloon and remove COOPDECH Endobronchial Blocker Tube.
- COOPDECH Endobronchial Blocker Tube can be used in conjunction with tubes intubated orotracheally, nasotracheally or by tracheotomy.
- The unique automatic inflation system enables the operator to inflate the cuff with one hand and operate bronchi fiberscope simultaneously.
- Joint Connector permits simultaneous sealed introduction of a bronchoscope and blocker while maintaining ventilation.
- Since the product is used in combination with a single-lumen tube and is used for blockage of the left or right lung procedure, there is no need to stock different sizes.
- This device may be connected to various types of tracheal tubes that include spiral types, as well as tracheotomy tubes and laryngeal masks.








COOPDECH Endobronchial Blocker Tube brings simplicity to endobronchial blockade of the right or left lung for procedures requiring low invasive one-lung ventilation.


Its required usage of single-lumen endotracheal tube eliminates potential loss of the airway, commonly posed during the extubation and re-intubation of a double-lumen tube.

Low-pressure, high-volume balloon creates excellent cuff surface contact with the inner bronchial wall, minimizing potential trauma to the bronchus.

## **COOPDECH Endobronchial Blocker Tube**


COOPDECH Endobronchial Blocker Tube is designed to achieve single-lung ventilation, and is used in combination with various other tubes inserted to secure the airways. The clinician can easily move the blocker from one lung to another. Manually torque the device on the shaft for precise placement into the desired airway. The angled tip let you easily choose sides for single-lung ventilation and provides a wide range of adjustment to precisely direct the blocker. Administration of oxygen to the collapsed lung, deaeration, and aspiration of secreted materials can be done via the suction port.





### **Auto Inflator**

Auto Inflator (Type B) can be operated with one hand if the balloon is inflated with air beforehand. Since the cuff can be inflated with a click of a button, operators can maneuver it by themselves without any help, while working on the bronchial fibers. Since the air is not injected directly via a syringe, damage to the bronchial tissues caused by high pressure and excessive injections can be minimized.



#### **Joint Connector**

COOPDECH Endbronchial Blocker Tube does not require re-intubation at the conclusion of surgery as with a double-lumen endotracheal tube. Joint connector can be used by connecting a bronchial blocker tube to various endotracheal tubes and anesthetic circuits. COOPDECH Endbronchial Blocker Tube is placed coaxially through a conventional endotracheal tube using a bronchoscope.



#### Cuff

Low-pressure barrel shaped balloon creates large cuff surface contact with the inner bronchial wall, minimizing potential trauma to the bronchus.

A soft light-blue colored cuff features a small outer diameter while deflated, preserving maximum airway volume for enhanced patient ventilation during thoracic procedures. Two types are available: rectangular round-shaped cuffs aimed at minimizing invasion, and small spindle-shaped cuffs that reduced airway resistance during cuff deaeration and realized even more enhanced visibility.

#### (Rectangular Round-shaped Cuffs)





(Small Spindle-shaped Cuffs)



